

The diagnosis and management of benzodiazepine dependence

Heather Ashton

Purpose of review

Despite repeated recommendations to limit benzodiazepines to short-term use (2–4 weeks), doctors worldwide are still prescribing them for months or years. This over-prescribing has resulted in large populations of long-term users who have become dependent on benzodiazepines and has also led to leakage of benzodiazepines into the illicit drug market. This review outlines the risks of long-term benzodiazepine use, gives guidelines on the management of benzodiazepine withdrawal and suggests ways in which dependence can be prevented.

Recent findings

Recent literature shows that benzodiazepines have all the characteristics of drugs of dependence and that they are inappropriately prescribed for many patients, including those with physical and psychiatric problems, elderly residents of care homes and those with comorbid alcohol and substance abuse. Many trials have investigated methods of benzodiazepine withdrawal, of which the keystones are gradual dosage tapering and psychological support when necessary. Several studies have shown that mental and physical health and cognitive performance improve after withdrawal, especially in elderly patients taking benzodiazepine hypnotics, who comprise a large proportion of the dependent population.

Summary

Benzodiazepine dependence could be prevented by adherence to recommendations for short-term prescribing (2–4 weeks only when possible). Withdrawal of benzodiazepines from dependent patients is feasible and need not be traumatic if judiciously, and often individually, managed.

Keywords

benzodiazepine dependence, benzodiazepine withdrawal, prevention of dependence

Abbreviation

GABA γ -aminobutyric acid

© 2005 Lippincott Williams & Wilkins
0951-7367

Introduction

Since their introduction in the 1950s, benzodiazepines appear to have passed their zenith of medical popularity. However, they are still prescribed excessively and often inappropriately. With their reputation perhaps approaching a nadir, at least as prescribed medications for long-term use, it is timely to review approaches to the diagnosis and management of dependence on these drugs.

The benzodiazepine bonanza

In the late 1970s benzodiazepines became the most commonly prescribed of all drugs in the world. Their range of actions – sedative/hypnotic, anxiolytic, anti-convulsant and muscle relaxant – combined with low toxicity and alleged lack of dependence potential seemed to make them ideal medications for many common conditions (Table 1). The drugs were prescribed long term, often for many years, for complaints such as anxiety, depression, insomnia and ordinary life stresses. Benzodiazepines were undoubtedly efficacious at first for these conditions, and apparently harmless – but there was a sting in the tail.

By the early 1980s long-term prescribed users themselves had realized that the drugs tended to lose their efficacy over time and instead became associated with adverse effects. In particular, patients found it difficult to stop taking benzodiazepines because of withdrawal reactions and many complained that they had become ‘addicted’ [1•]. Controlled clinical trials among such patients [2–4] demonstrated beyond doubt that withdrawal symptoms, even from regular ‘therapeutic’ doses of benzodiazepines, were real and that they indicated dependence on the drugs.

Changing definitions of dependence

That benzodiazepines could cause physical dependence was accepted by the medical profession on the basis that a withdrawal syndrome occurred on cessation of regular use, and doctors were advised to reserve them for short-term use in minimal dosage [5,6]. However, definitions of drug dependence changed in the 1990s. Previously, dependence had been defined in terms of

Curr Opin Psychiatry 18:249–255. © 2005 Lippincott Williams & Wilkins.

Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne, UK

Correspondence to Professor C.H. Ashton, Department of Psychiatry, University of Newcastle upon Tyne, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP, UK
Tel: +44 191 2226000 ext 6978; fax: +44 191 2226162;
e-mail: c.h.ashton@ncl.ac.uk

Current Opinion in Psychiatry 2005, 18:249–255

Table 1. Therapeutic actions of benzodiazepines (in short-term use)

Action	Clinical use
Anxiolytic – relief of anxiety	Anxiety and panic disorders, phobias Agitated psychoses
Hypnotic – promotion of sleep	Insomnia
Myorelaxant – muscle relaxation	Muscle spasms, spastic disorders
Anticonvulsant – stops fits, convulsions	Fits due to drug poisoning, some forms of epilepsy, alcohol withdrawal
Amnesia – impairment of short-term memory	Premedication for operations, sedation for minor surgical operations

the development of drug tolerance and a withdrawal syndrome on cessation, but in current classification systems these two features alone are no longer considered sufficient for the diagnosis. Present criteria for substance dependence [7] include tolerance, escalation of dosage, continued use despite efforts to stop and knowledge of adverse effects, other behavioural features, and a withdrawal syndrome (Table 2). Benzodiazepines meet all these criteria.

Tolerance and dosage escalation

Tolerance to benzodiazepines develops at different rates and to different degrees for the various actions. Tolerance to hypnotic effects develops rapidly, within a few days or weeks of regular use. Studies in elderly patients indicate that, when taken over long periods, benzodiazepines have little effect on sleep [8,9^{••},10[•]]. Although some poor sleepers report continued efficacy of benzodiazepine hypnotics, possibly because they prevent rebound insomnia (a withdrawal effect), clinical experience shows that a considerable proportion of hypnotic users gradually increase their dosage, sometimes to above recommended levels. It is not uncommon for insomniacs to be taking two or more nightly benzodiazepines concurrently [11^{••},12[•]].

Tolerance to the anxiolytic effects of benzodiazepines develops more slowly, over a few months, and clinical

observations show that long-term use does little to control, and may even aggravate, anxiety [13]. There is also evidence of dosage escalation in anxiolytic users. In one clinical study over 25% of the patients were taking two benzodiazepines, the second having been added to the prescription when the first ceased to be effective [13]. Although some authors recommend long-term use of benzodiazepine anxiolytics for certain conditions [14,15], it is likely that the drugs are preventing withdrawal symptoms rather than reducing anxiety [16].

Tolerance to the anticonvulsant effects of benzodiazepines occurs within a few weeks in a high proportion of patients with epilepsy [17] and also to the muscular relaxant effects when used in patients with spastic disorders. Of particular clinical importance, however, is the finding that little tolerance develops to the amnesic effects and other cognitive impairments caused by benzodiazepines. Studies of long-term users have shown deficits in learning, memory, attention and visuospatial ability. A metaanalysis of 13 research studies revealed moderate–large deficits in all 12 of the cognitive domains tested in long-term benzodiazepine users compared with controls [18^{••}]. Such effects are most marked in the elderly in whom they may suggest dementia [19]. Improvement occurs when the drugs are stopped, but it may be slow and perhaps incomplete [20^{••},21].

Escalation of dosage and chronic use of benzodiazepines cause additional adverse effects including depression, excessive sedation, leading to falls and fractures, road traffic and other accidents (especially when combined with alcohol), and the insidious development of increasing psychological and physical symptoms [13,16,21,22, 23[•]–25[•]]. Again, the elderly are most vulnerable to these effects, especially if taking multiple medications [26^{••}]. Furthermore, benzodiazepines can be lethal in overdose [27,28[•]].

Withdrawal syndrome

The existence of a benzodiazepine withdrawal syndrome has been abundantly demonstrated [2–4,29,30]. The

Table 2. Criteria for substance dependence^a

1	Tolerance as defined by either a need for markedly increased amounts of the substance to achieve the clinical effect, or markedly diminished effect with continued use of the same amount of the substance
2	Withdrawal as defined by either the characteristic withdrawal syndrome for the substance, or the same or similar substance is taken to avoid withdrawal symptoms
3	The substance is taken in larger amounts or over a longer period than was intended
4	There is a persistent desire or unsuccessful attempts to cut down or control substance use
5	Time is spent in activities necessary to obtain the substance (e.g. visiting multiple doctors)
6	Important activities are given up or reduced because of substance use
7	The substance use is continued despite knowledge of having a problem caused or exacerbated by the substance

^aA maladaptive pattern of substance use, leading to clinically significant impairment or distress as manifested by three or more of the above, occurring at any time in the same 12-month period. Reprinted with permission from the Diagnostic and Statistical Manual of Mental Disorders, copyright 2000. American Psychiatric Association [7].

Table 3. Some common benzodiazepine withdrawal symptoms

Symptoms common to all anxiety states	Symptoms less common in anxiety states – relatively specific to benzodiazepine withdrawal ^a
Anxiety, panic attacks, agoraphobia	Perceptual distortions, sense of movement
Insomnia, nightmares	Depersonalization, derealization
Depression, dysphoria	Hallucinations (visual, auditory)
Excitability, restlessness	Distortion of body image
Poor memory and concentration	Tingling, numbness, altered sensation
Dizziness, light headedness	Formication (skin 'crawling')
Weakness 'jelly legs'	Sensory hypersensitivity (light, sound, taste, smell)
Tremor	Muscle twitches, jerks, fasciculation
Muscle pain, stiffness	Tinnitus
Sweating, night sweats	Psychotic symptoms ^a
Palpitations	Confusion, delirium ^a
Blurred or double vision	Convulsions ^a

^aUsually only on rapid or abrupt withdrawal from high doses of benzodiazepines.

syndrome can be mild and short-lived or severe and sometimes protracted [31]. Symptoms include many that are common to anxiety states in general, as well as some more characteristic of benzodiazepine withdrawal (Table 3). Severity is often associated with prolonged or high-dose use, short-acting potent benzodiazepines, certain personality types and anxiety/neuroticism [32,33]. The reported incidence varies between 30 and 100% in different studies, but up to 50% of long-term users decline to participate in, or drop out of, withdrawal studies [4,34,35]. Withdrawal symptoms prolong benzodiazepine use, which often continues for years after the initial indication for the drug has passed. Many long-term users, aware that the drugs are no longer effective or are causing adverse effects, have tried to stop but have been unsuccessful because of the emergence of withdrawal symptoms [13].

Mechanisms of tolerance and withdrawal

The pharmacological mechanisms underlying benzodiazepine tolerance and withdrawal are complex and still not clear. Present knowledge has recently been reviewed in detail [36,37]. Tolerance to chronic benzodiazepine administration appears to result from neuroadaptive processes involving both desensitization of inhibitory γ -aminobutyric acid (GABA) receptors and sensitization of excitatory glutamergic receptors. Both these systems include multiple receptor subtypes. Changes in GABA receptors may include conformational alterations towards a low affinity state for GABA and uncoupling of benzodiazepine receptors from their sites on certain GABA_A receptors, followed by internalization and perhaps long-term effects on intraneural gene transcription [36,38^{*}]. Changes in the glutamergic system may include sensitization of *N*-methyl-D-aspartate (NMDA) and possibly other receptors

[1^{**},37]. These adaptations could occur on different time scales depending on the receptor subtype and brain region involved, thus accounting for the differing rates of development of tolerance to various benzodiazepine actions.

Rapid or abrupt withdrawal of the benzodiazepine once tolerance has developed exposes the recipient to the consequences of all these drug-induced adaptations. The result is underactivity of inhibitory GABA functions and a surge in excitatory nervous activity, giving rise to many of the benzodiazepine withdrawal symptoms shown in Table 3. The various receptor changes occurring during tolerance may be slow to reverse and may do so at different rates, possibly explaining the variable time of emergence and duration of individual withdrawal symptoms and sometimes protracted nature of benzodiazepine withdrawal [31].

Diagnosis of benzodiazepine dependence

The key signs of benzodiazepine dependence are withdrawal symptoms on dosage reduction or discontinuation. However, dependence can often be inferred in continuing benzodiazepine users from a history of long-term use, reliance on regular prescriptions, dosage escalation, unsuccessful attempts to cut down or stop drug use, and high anxiety levels [39^{*},40^{*},41,42^{**},43]. Chronic benzodiazepine users with a history of other drug or alcohol dependence are also likely to be dependent [44]. Unfortunately, long-term prescription of benzodiazepines continues today [45^{*}] and maintains several overlapping populations of benzodiazepine-dependent users.

Benzodiazepine-dependent populations

There are three overlapping types of benzodiazepine-dependent populations.

Therapeutic dose dependence

The largest population of benzodiazepine-dependent patients comprises long-term users who have inadvertently become dependent as a result of regular repeat prescriptions over months or years. The size of this population is estimated as 500 000 to 1 million in the UK, 4 million in the US [1^{**}] and several million worldwide [46]. It is likely that at least 50% of these users are dependent. A considerable proportion of these patients are elderly females taking benzodiazepine hypnotics [8,9^{**}] and it is noteworthy that prescriptions for hypnotics (including drugs with similar actions such as zopiclone) have not declined despite a reduction in prescriptions for benzodiazepine anxiolytics [39^{*}]. Other long-term prescribed users who are likely to be dependent are patients with physical and psychiatric problems and elderly residents of care homes [46,47^{*},48^{*},49,50^{*}].

Prescribed high-dose dependence

A minority of patients who start on prescribed benzodiazepines escalate their dosage excessively. At first they may persuade their doctors to increase prescriptions, but on reaching the prescriber's limits, they may attend several doctors or hospital departments to obtain further supplies. When other sources fail they may resort to 'street' benzodiazepines.

Recreational benzodiazepine abuse

The use of benzodiazepines as recreational drugs is a growing problem [16]. The size of this population is unknown but estimates suggest about 200 000 people in the UK alone (population 55 million) and similar or higher proportions in the US, Europe, Australia and other countries. Benzodiazepines commonly form part of a polysubstance abuse pattern. They are taken by at least half of opiate, amphetamine, cocaine and other illegal drug users worldwide and by alcoholics [1^{••},51]. Other users include patients with mental illness and comorbid other substance abuse [48[•],49]. Some people use benzodiazepines as their primary recreational drug, bingeing intermittently on high doses or injecting intravenously with risk of gangrene, HIV and hepatitis C [51].

Reasons given for taking benzodiazepines recreationally are that they enhance the 'high' obtained from illicit drugs, alleviate withdrawal effects, serve as 'downers' from the effects of stimulant drugs ('uppers') and also produce a 'kick' when taken alone in high doses or injected intravenously. Many illicit benzodiazepine users become dependent and show typical withdrawal symptoms which can be severe [52,53]. The tragedy of recreational benzodiazepine abuse is that it is largely iatrogenic, resulting from widespread overprescription of benzodiazepines which increased their general availability. Major sources of illicit benzodiazepines are from general practitioner prescriptions and thefts from chemists or pharmaceutical warehouses [51]. They are available on the black market [54[•]] and can be obtained on the Internet.

Management of benzodiazepine withdrawal

Because of the adverse effects, lack of efficacy and socio-economic costs of continued benzodiazepine use, long-term users have for many years been advised to withdraw if possible or at least to reduce dosage [5,45[•]]. However, benzodiazepine withdrawal has often been badly managed and has acquired a reputation as a traumatic process for both patients and doctors. This reputation is largely undeserved if the process is carried out judiciously. The management of withdrawal has been reviewed by many authors [1^{••},55–57,58[•]]. All agree that the key strategies for successful discontinuation are gradual dosage tapering and psychological support if necessary.

Dosage reduction

The rate of dosage reduction varies for different types of benzodiazepine patients.

Therapeutic dose users

Benzodiazepine dosage should be tapered gradually since abrupt withdrawal, especially from high doses, can precipitate convulsions, acute psychotic states and other severe reactions (Table 3). The recommended rate of tapering for patients on therapeutic doses of benzodiazepine is withdrawal in steps of about one-eighth to one-tenth of the daily dose every 1–2 weeks [59[•],60]. Over-rapid tapering such as fixed dosage reductions of 25–50% every 1 or 2 weeks or faster, especially in patients taking potent or rapidly eliminated benzodiazepines (Table 4), increases the likelihood of other withdrawal symptoms, dropouts from trials, need for psychological support and later relapse [11^{••},12[•],35]. For this reason, the rate of withdrawal should be individually adjusted to the patient's needs, taking into account factors such as dosage and type of benzodiazepine, reasons for prescription, lifestyle, personality, environmental stresses and amount of available support. Various authors have suggested optimal times of 6–8 weeks for withdrawal but some patients may require a year or more [1^{••},59[•]]. Ideally, after receiving advice and information from the physician and giving full consent, the patient should be in control of his/her own personal reduction rate and proceed at whatever pace is tolerable. A personalized approach is likely to result in fewer patients dropping out or declining to participate in withdrawal trials.

In general practice settings, even minimal intervention such as a letter with an information sheet or a single brief consultation can be effective in reducing or stopping benzodiazepine use without adverse effects. In one

Table 4. Approximate equivalent doses and elimination half-lives of benzodiazepines

Benzodiazepine	Approximately equivalent dosage (mg) ^a	Elimination half-life (h) (active metabolite)
Alprazolam	0.5	6–12
Chlordiazepoxide	25	5–30 (36–200)
Clonazepam	0.5	18–50
Diazepam	10	20–100 (36–200)
Flunitrazepam	1	18–26 (36–200)
Flurazepam	15–30	(40–250)
Loprazolam	1	6–12
Lorazepam	1	10–20
Lormetazepam	1	10–12
Nitrazepam	10	15–38
Oxazepam	20	4–15
Temazepam	20	8–22

^aClinical potency for hypnotic or anxiolytic effects may vary between individuals; equivalent doses are approximate.

controlled study of 191 mainly elderly long-term hypnotic users, within 6 months such measures resulted in significant dosage reduction or complete cessation [9^{••}]. The advice in the letter sent to patients was simply 'try reducing by half a tablet every few weeks'. Those who reduced dosage by 25% or more showed improvement in mental and physical health, reported no withdrawal symptoms or sleep problems, and required fewer medical consultations. In another controlled general practice study, 192 elderly hypnotic users underwent a tapered dosage programme over 8–9 weeks using placebo [8]. Eighty percent had successfully withdrawn by 6 months. These patients showed improvements in cognitive and psychomotor performance and did not differ in sleep or withdrawal symptoms from a control group who continued taking benzodiazepines.

For some patients, particularly those taking benzodiazepines for anxiety or using potent benzodiazepines (lorazepam, alprazolam, clonazepam) (Table 4), there are advantages in conducting the withdrawal by using diazepam. The slow elimination of this drug ensures a gradual fall in blood concentration while its availability in low-dosage forms permits small dosage reductions. Conversion from other benzodiazepines to diazepam can be conducted in stages, and it is important to allow for equivalent potencies between different benzodiazepines (Table 4). Full details and examples of withdrawal schedules from different benzodiazepines at various doses are available on the Internet [57] and other references [1^{••},56,60].

High-dose abusers

A different withdrawal approach is required for high-dose benzodiazepine abusers in whom benzodiazepine use often forms a part of polydrug abuse pattern. These patients may need in-patient detoxification for the primary drug and a fairly rapid withdrawal of the benzodiazepine, with diazepam substitution and tapering over 2–3 weeks being usual [52,53]. The development of convulsions can usually be prevented by moderate doses of diazepam (10 mg), but some authors report benefit from carbamazepine [61,62].

Adjuvant drugs

Many drugs have been investigated for their ability to attenuate benzodiazepine withdrawal symptoms, but none has proved generally useful for patients dependent on therapeutic doses [1^{••}]. Drugs tested include antidepressants [34,63], β -blockers [4], buspirone [64,65], carbamazepine and other anticonvulsants [61,62,66[•]], flumazenil [1^{••},67,68], captodiamine [69[•]], gabapentin [70] and others are under investigation [71,72[•],73[•]]. Nonbenzodiazepine GABA receptor agonists such as zopiclone relieve withdrawal symptoms but are contra-

indicated since they have the same disadvantages as benzodiazepines [74[•]].

Psychological support

The degree of psychological support required during withdrawal is variable and may range from a single brief consultation or letter [9^{••}] to more formal cognitive, behavioural or other therapies directed towards anxiety management and stress-coping strategies [1^{••},56]. Support when needed should be available both during and after withdrawal since patients may remain vulnerable to stress for some months. Information about withdrawal symptoms should be supplied and referral to a support organization is often helpful.

Outcome of withdrawal

With carefully managed withdrawal and adequate psychological support in motivated patients, the success rate for stopping benzodiazepines can be 70–80% [8,13,34,35]. Successful cessation need not be affected by duration of use, type or dosage of benzodiazepines, severity of symptoms, psychiatric history or personality disorder, although symptom severity is greater in anxious individuals [13,32,33,65]. Relapse rates 1–5 years after withdrawal vary between 8 and 57% in different studies [8,11^{••},13,34,75] but are probably minimized by using individualized withdrawal programmes. Some patients revert temporarily to benzodiazepine use after withdrawal but most stop again or considerably reduce dosage [34,75].

Conclusion

Prevention of benzodiazepine dependence can be achieved by adherence to official recommendations to limit prescriptions to short-term use (2–4 weeks), or as intermittent brief courses or occasional doses [5,6]. Although prescribing of benzodiazepines has declined substantially since 1988, 30% of GP prescriptions in the UK are still for 56 or more tablets [45[•]] and many physicians in Europe, the US and Australia continue to prescribe them long term. Particular care should be taken in prescribing benzodiazepines for vulnerable patients such as those with alcohol or drug dependence, and doctors should be aware that prescriptions may enter the illicit market. Benzodiazepines are not indicated for long-term treatment of depression and when used for chronic psychiatric conditions such as schizophrenia, bipolar affective disorder, anxiety disorders and chronic insomnia, clinicians should examine the risk–benefit ratio at an early stage so that the risks of dependence can be balanced against any therapeutic benefits [76–79,80[•]]. Finally, doctors should avoid using nonbenzodiazepine hypnotics and anxiolytics such as zopiclone, zolpidem and zaleplon in benzodiazepine-dependent patients since these drugs can also cause dependence and abuse.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

- 1 Ashton H. Benzodiazepine dependence. In: Haddad P, Dursun S, Deakin B, editors. *Adverse syndromes and psychiatric drugs*. Oxford: Oxford University Press; 2004. pp. 239–260.
- A full review of benzodiazepine dependence including its recognition, descriptions of the clinical features and pharmacological mechanisms of tolerance and withdrawal reactions, benzodiazepine-dependent populations, management of withdrawal, and prevention of dependence.
- 2 Hallstrom C, Lader MH. Benzodiazepine withdrawal phenomena. *Int Pharmacopsychiatry* 1981; 16:235–244.
- 3 Petursson H, Lader MH. Benzodiazepine dependence. *Br J Addict* 1984; 76:133–145.
- 4 Tyrer P, Rutherford D, Higgitt T. Benzodiazepine withdrawal symptoms and propranolol. *Lancet* 1981; 1:520–522.
- 5 Committee on Safety of Medicines. Benzodiazepines, dependence and withdrawal symptoms. *Current Problems* 21; 1988.
- 6 Royal College of Psychiatrists. Benzodiazepines and dependence: a college statement. *Bull R Coll Psychiatrists* 1988; 12:107–108.
- 7 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders (DSM-IV)*. Washington, DC: APA; 1994.
- 8 Curran HV, Collins R, Fletcher S, et al. Older adults and withdrawal from benzodiazepine hypnotics in general practice: effects on cognitive function, sleep, mood and quality of life. *Psychosom Med* 2003; 33:1223–1237.
- 9 Heather N, Bowie A, Ashton H, et al. Randomised controlled trial of two brief interventions against long-term benzodiazepine use: outcome of intervention. *Addict Res Theory* 2004; 12:141–154.
- A controlled UK general practice trial describing the results of brief interventions (letter or single short consultation) on benzodiazepine consumption in 273 long-term prescribed users, mainly elderly patients taking benzodiazepine hypnotics.
- 10 Holbrook AM. Treating insomnia [editorial]. *BMJ* 2004; 329:1198–1199.
- Editorial discussing the increasing use of hypnotic drugs despite little evidence of clinical benefit. Hypnotics described as examples of 'iatrogenesis imperfecta'.
- 11 Morin CM, Belanger L, Bastien C, Vallieres A. Long-term outcome after discontinuation of benzodiazepines for insomnia: a survival analysis of relapse. *Behav Res Ther* 2005; 43:1–14.
- A Canadian study investigating medication-free survival time 2 years after tapered discontinuation and cognitive-behavioural therapy in older patients with insomnia. A substantial relapse rate was found, although this was less in those who received cognitive-behavioural therapy.
- 12 Poyares D, Guilleminault C, Ohayon C, Tufik S. Chronic benzodiazepine usage and withdrawal in insomnia patients. *J Psychiatr Res* 2004; 38:327–334.
- Sleep quality including slow wave sleep improved after benzodiazepine discontinuation in 25 Brazilian patients with persistent insomnia.
- 13 Ashton H. Benzodiazepine withdrawal: outcome in 50 patients. *Br J Addict* 1987; 83:665–671.
- 14 Hollister LE, Muller-Oerlinghausen B, Rickels K, et al. Clinical uses of benzodiazepines. *J Clin Psychopharmacol* 1993; 13(Suppl 1):1S–169S.
- 15 Schweizer E, Rickels K, Weiss S, Zavodnick S. Maintenance drug treatment of panic disorder. I: Results of a prospective, placebo-controlled comparison of alprazolam and imipramine. *Arch Gen Psychiatry* 1993; 50:51–60.
- 16 Griffiths RR, Weerts EM. Benzodiazepine self-administration in humans and laboratory animals: implications for problems of long-term use and abuse. *Psychopharmacology* 1997; 134:1–37.
- 17 Brodie MJ. Established anticonvulsants and treatment of refractory epilepsy. *BMJ* 1998; 336:350–354.
- 18 Barker MJ, Greenwood KM, Jackson M, Crowe SF. Cognitive effects of long-term benzodiazepine use: a meta-analysis. *CNS Drugs* 2004; 18: 37–48.
- A metaanalysis of 13 research investigations into the effects of long-term benzodiazepine use on cognition. Long-term benzodiazepine users were significantly impaired compared with controls across all 12 of the cognitive domains tested.
- 19 Lader M. Benzos and memory loss: more than just 'old age'. *Prescriber* 1992; 3:13.
- 20 Barker MJ, Greenwood KM, Jackson M, Crowe SF. Persistence of cognitive effects after withdrawal from long-term benzodiazepine use: a meta-analysis. *Arch Clin Neuropsychol* 2004; 19:437–454.
- A metaanalysis of cognitive functioning in long-term benzodiazepine users after withdrawal, showing some recovery of function but a significant remaining impairment in most areas of cognition compared with controls, at least 6 months after cessation. The results suggest that there may be some permanent deficits.
- 21 Gray SL, Penninx BW, Blough DK, et al. Benzodiazepine use and physical performance in community dwelling older women. *J Am Geriatr Soc* 2003; 51:1563–1570.
- 22 Panneman MJ, Goettsch WG, Kramarz P, Herings RM. The costs of benzodiazepine-associated hospital-treated fall injuries in the EU: a Pharmo study. *Drugs Aging* 2003; 20:833–839.
- 23 Wagner AK, Zhang F, Soumerai SB, et al. Benzodiazepine use and hip fractures in the elderly: who is at greatest risk? *Arch Intern Med* 2004; 164:1567–1572.
- Examination of the association between benzodiazepine use and hip fractures in a US population.
- 24 Walsh JM, Flegel R, Cangianelli LA, et al. Epidemiology of alcohol and other drug use among motor vehicle crash victims admitted to a trauma center. *Traffic Inj Prev* 2004; 5:254–260.
- A US study of the incidence and prevalence of alcohol and other drug use in patients injured in vehicle crashes. Benzodiazepines were the most frequently detected drugs after alcohol and marijuana.
- 25 Movig KL, Mathijssen MP, Nagel PH, et al. Psychoactive substance use and the risk of motor vehicle accidents. *Accid Anal Prev* 2004; 36:631–636.
- Study of vehicle accidents requiring hospitalization in the Netherlands, showing that single use of benzodiazepines as well as combinations with alcohol and other drugs increases the risks.
- 26 Madhusoodanan S, Bogunovic OJ. Safety of benzodiazepines in the geriatric population. *Expert Opin Drug Safety* 2004; 3:485–493.
- A comprehensive description of the special risks of benzodiazepines in the elderly, with description of age-related pharmacokinetic and pharmacodynamic factors.
- 27 Stenhouse G, Grieve JH. Drug-related deaths in Grampian, Scotland. *Scott Med J* 2003; 48:111–113.
- 28 Isbister GK, O'Regan L, Sibbritt D, Whyte IM. Alprazolam is relatively more toxic than other benzodiazepines in overdose. *Br J Clin Pharmacol* 2004; 58:88–95.
- Study from Australia showing that alprazolam was more toxic in overdose due to self-poisoning than other benzodiazepines.
- 29 Tyrer P, Owen R, Dawling S. Gradual withdrawal of diazepam after long-term therapy. *Lancet* 1983; 1:1402–1406.
- 30 Owen RT, Tyrer P. Benzodiazepine dependence: a review of the evidence. *Drugs* 1983; 25:385–398.
- 31 Ashton H. Protracted withdrawal from benzodiazepines: the post-withdrawal syndrome. *Psychiatr Ann* 1995; 25:158–165.
- 32 Murphy SM, Tyrer P. The essence of benzodiazepine dependence. In: Lader M, editor. *The psychopharmacology of addiction*. Oxford: Oxford University Press; 1991. pp. 157–167.
- 33 O'Connor KP, Marchand A, Belanger L, et al. Psychological distress and adaptational problems associated with benzodiazepine withdrawal and outcome: a replication. *Addict Behav* 2004; 2:583–593.
- 34 Zitman FG, Couvee JE. Chronic benzodiazepine use in general practice patients with depression: an evaluation of controlled treatment and taper-off. *Br J Psychiatry* 2001; 178:317–324.
- 35 Baillargeon L, Landreville P, Verreault R, et al. Discontinuation of benzodiazepines among older insomniac adults treated with cognitive-behavioural therapy combined with gradual tapering: a randomized trial. *CMAJ* 2003; 169:1015–1020.
- 36 Bateson AN. Basic pharmacologic mechanisms involved in benzodiazepine tolerance and withdrawal. *Curr Pharmaceut Des* 2002; 8:5–21.
- 37 Allison C, Pratt JA. Neuroadaptive processes in GABAergic and glutamatergic systems in benzodiazepine dependence. *Pharmacol Ther* 2003; 98:171–195.
- 38 Huopaniemi L, Keist R, Randolph A, et al. Diazepam-induced adaptive plasticity revealed by alpha1 GABA_A receptor-specific expression profiling. *J Neurochem* 2004; 88:1059–1067.
- Study of down-regulation of GABA_A receptors following diazepam administration in mutated mice relevant to mechanisms of benzodiazepine tolerance and dependence.
- 39 What's wrong with prescribing hypnotics? *Drug Ther Bull* 2005; 42:89–93.
- A review of the risks associated with long-term hypnotic use and how effects can be minimized.

- 40 Voyer P, McCubbin M, Cohen D, *et al.* Unconventional indicators of drug dependence among elderly long-term users of benzodiazepines. *Issues Ment Health Nurs* 2004; 25:603–628.
A study of users' reports of factors indicating dependence in 45 elderly persons taking long-term benzodiazepines.
- 41 Morin CM, Belanger L, Bernier F. Correlates of benzodiazepine use in individuals with insomnia. *Sleep Med* 2004; 5:457–462.
- 42 Kan CC, Hilberink SR, Breteler MH. Determination of the main risk factors for benzodiazepine dependence using a multivariate and multidimensional approach. *Compr Psychiatry* 2004; 45:88–94.
A careful study from the Netherlands using regression analyses and validated screening questionnaires in 599 outpatients to determine predictors of benzodiazepine dependence severity.
- 43 Neutel CI, Walop W, Patten SB. Can continuing benzodiazepine use be predicted? *Can J Clin Pharmacol* 2003; 10:202–206.
- 44 Johansson BA, Berglund M, Hanson M, *et al.* Dependence on legal psychotropic drugs among alcoholics. *Alcohol* 2003; 38:613–618.
- 45 CMO's Update 37. Benzodiazepines warning. Department of Health; January 2004. p. 4.
Reminder to UK doctors that benzodiazepines should be used for 2–4 weeks only, but 30% of GP prescriptions were for 56 or more tablets.
- 46 Ashton H, Golding JF. Tranquillisers: prevalence, predictors and possible consequences. Data from a large United Kingdom survey. *Br J Addict* 1989; 84:541–546.
- 47 Valenstein M, Taylor KK, Austin K, *et al.* Benzodiazepine use among depressed patients treated in mental health settings. *Am J Psychiatry* 2004; 161:654–661.
A study of inappropriate benzodiazepine prescribing among depressed patients in mental health settings in the US, involving 128 029 patients identified from the Veterans Affairs National Registry.
- 48 Clark RE, Xie H, Brunette MF. Benzodiazepine prescription practices and substance abuse in persons with severe mental illness. *J Clin Psychiatry* 2004; 65:151–155.
Study from the US showing greater benzodiazepine use in patients with severe mental illness and comorbid substance use than in patients with mental disorder alone.
- 49 Brunette MF, Noordsy DL, Xie H, Drake RE. Benzodiazepine use and abuse among patients with severe mental illness and co-occurring substance use disorders. *Psychiatr Serv* 2003; 54:1395–1401.
- 50 Zandstra SM, Van Rijswijk E, Rijnders CA, *et al.* Long-term benzodiazepine users in family practice: differences from short-term users in mental health, coping behaviour and psychological characteristics. *Fam Pract* 2004; 21:266–269.
Dutch study of patient characteristics related to long-term benzodiazepine use in general practice.
- 51 Ashton CH. Benzodiazepine abuse. In: Caan E, de Bellerocche J, editors. *Drink, drugs and dependence*. London: Routledge; 2002. pp. 197–212.
- 52 Seivewright N, Dougal W. Withdrawal symptoms from high dose benzodiazepines in polydrug users. *Drug Alcohol Depend* 1993; 32:15–23.
- 53 Seivewright N. Theory and practice in managing benzodiazepine dependence and misuse. *J Subst Misuse* 1998; 3:170–177.
- 54 Grzybowski S. The black market in prescription drugs. *Med Crime Punishment* 2004; 364:28–29.
Black market entrepreneurship of prescription drugs in developed and developing countries.
- 55 Lader M. Avoiding long-term use of benzodiazepine drugs. *Prescriber* 1991; March:79–93.
- 56 Ashton H. The treatment of benzodiazepine dependence. *Addiction* 1994; 89:1535–1541.
- 57 Ashton CH. Benzodiazepines: how they work and how to withdraw; 2002. <http://www.benzo.org.uk>. [Accessed February 2005].
- 58 Lingford-Hughes AR, Welch S, Nutt DJ. Evidence-based guidelines for the pharmacological management of substance misuse, addiction and comorbidity: recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2004; 18:293–335.
Guidelines and recommendations for the management of various drug dependencies including benzodiazepine misuse and dependence.
- 59 British National Formulary. British Medical Association and Royal Pharmaceutical Society of Great Britain; 2004.
Official advice on benzodiazepine withdrawal and the use of diazepam substitution.
- 60 Ashton H. How to wean patients off benzodiazepines. *Pulse* 2004; 64:50–53.
- 61 Ries RK, Roy-Birynne PP, Ward NG, *et al.* Carbamazepine treatment for benzodiazepine withdrawal. *Am J Psychiatry* 1989; 146:536–537.
- 62 Schweizer E, Rickels K, Case WG, *et al.* Carbamazepine treatment in patients discontinuing long-term benzodiazepine therapy: effects on withdrawal severity and outcome. *Arch Gen Psychiatry* 1991; 48:448–452.
- 63 Tyrer P, Ferguson B, Hallstrom C, *et al.* A controlled trial of dothiepin and placebo in treating benzodiazepine withdrawal symptoms. *Br J Psychiatry* 1996; 168:457–461.
- 64 Olajide D, Lader M. A comparative study of the efficacy of buspirone in relieving benzodiazepine withdrawal symptoms. *J Clin Psychopharmacol* 1987; 7:11–15.
- 65 Ashton CH, Rawlins MD, Tyrer SP. A double-blind placebo-controlled study of buspirone in diazepam withdrawal in chronic benzodiazepine users. *Br J Psychiatry* 1990; 157:232–238.
- 66 Zullino DF, Khazaal Y, Hattenschwiler J, *et al.* Anticonvulsant drugs in the treatment of substance withdrawal. *Drugs Today (Barc)* 2004; 40:603–619.
Useful article discussing the potential of anticonvulsant drugs in alleviating withdrawal symptoms in substance abuse patients.
- 67 Gerra G, Zaimovic A, Giusti F, *et al.* Intravenous flumazenil versus oxazepam tapering in the treatment of benzodiazepine withdrawal: a randomized, placebo-controlled study. *Addict Biol* 2002; 7:385–395.
- 68 Savic I, Widen L, Stone-Elander S. Feasibility of reversing benzodiazepine tolerance with flumazenil. *Lancet* 1991; 337:133–137.
- 69 Mercier-Guyon C, Chabannes JP, Saviuc P. The role of captodiamine in the withdrawal from long-term benzodiazepine treatment. *Curr Med Res Opin* 2004; 20:1347–1355.
French trial of captodiamine in benzodiazepine discontinuation. This drug apparently alleviated withdrawal symptoms, but further studies are indicated.
- 70 Crockford D, White WD, Campbell B. Gabapentin use in benzodiazepine dependence and detoxification [abstract]. *Can J Psychiatry* 2001; 46:287.
- 71 Nemeroff CB. The role of GABA in the pathophysiology and treatment of anxiety disorders. *Psychopharmacol Bull* 2003; 37:136–146.
- 72 Basile AS, Lippa AS, Skolnick P. Anxiolytic effects: can less be more? *Eur J Pharmacol* 2004; 500:441–451.
A review of preclinical and clinical studies of selective anxiolytics which might limit benzodiazepine side effects.
- 73 Bateson AN. The benzodiazepine site of the GABA_A receptor: an old target with new potential? *Sleep Med* 2004; 5(Suppl 1):S9–S15.
A discussion of the pharmacological actions of benzodiazepines and other drugs acting at the same sites on GABA_A receptors.
- 74 Zaleplon, zolpidem and zopiclone for the short-term management of insomnia. National Institute for Clinical Excellence Technology Appraisal Guidance 77; 2004.
Cautions UK doctors concerning use of zaleplon, zolpidem and zopiclone in preference to benzodiazepines for short-term treatment of insomnia.
- 75 Holton A, Tyrer P. Five year outcome of patients withdrawn from long-term treatment with diazepam. *BMJ* 1990; 300:1241–1242.
- 76 Royal College of Psychiatrists. Benzodiazepines: risks, benefits and dependence: a re-evaluation. CR59; 2002.
- 77 Allgulander C, Bandelow B, Hollander E, *et al.* WCA recommendations for the long-term treatment of generalized anxiety disorder. *CNS Spectr* 2003; 9(Suppl 1):53–61.
- 78 Van Ameringen M, Allgulander C, Bandelow B, *et al.* WCA recommendations for the long-term treatment of social phobia. *CNS Spectr* 2003; 8(Suppl 1):40–52.
- 79 Pollack MH, Allgulander C, Bandelow B, *et al.* WCA recommendations for the long-term treatment of panic disorder. *CNS Spectr* 2003; 8(Suppl 1):17–30.
- 80 Jindal RD, Buysse DJ, Thase ME. Maintenance treatment of insomnia: what can we learn from the depression literature? *Am J Psychiatry* 2004; 161:19–24.
An overview of the management of insomnia, including a discussion on the advantages and disadvantages of long-term use of benzodiazepines and other hypnotic drugs.